

ACC-i2 with TCT

VOLUMETRIC ASSESSMENT OF CORONARY CALCIFICATION USING OPTICAL COHERENCE TOMOGRAPHY

i2 Poster Contributions

McCormick Place South, Hall A

Saturday, March 24, 2012, 9:30 a.m.-Noon

Session Title: Intravascular Diagnostics

Abstract Category: 3. Intravascular Diagnostics

Presentation Number: 2527-482

Authors: *Emile Mehanna, David Prabhu, Eric Brandt, Hirosada Yamamoto, Daniel Chamie, Guilherme Attizzani, Yusuke Fujino, Satoko Tahara, Nienke Van Ditzhuijzen, Wei Wang, Madhusudhana Gargesh, Hiram Bezerra, David Wilson, Marco Costa, Harrington-McLaughlin Heart and Vascular Institute, University Hospitals Case Medical Center, Cleveland, OH, USA, Department of Biomedical Engineering, Case Western Reserve University, Cleveland, OH, USA*

Background: Calcified plaques (CP) present unique challenges for percutaneous coronary intervention. We sought to quantify calcium volume in coronary atherosclerotic plaques by means of intravascular optical coherence tomography (OCT).

Methods: Thirty cadaveric coronary arteries were imaged using Frequency-Domain OCT. Vessels were frozen, automatically sectioned and imaged at 20 μ m intervals using an in-house built cryo-imaging system. Volumetric co-registration between OCT and cryo-images was performed. CP area, distance from lumen and angle were traced (fig 1) and total volume was calculated.

Results: Quantification was performed in 257 OCT and 1285 co-registered cryo-images (19 matched CP). Interpolation was required to trace deep calcification boundaries in 7 cases. Overall there was good agreement for measurement of luminal depth of calcium (0.25 ± 0.09 vs. 0.26 ± 0.12 mm), circumferential distribution of calcification (35.33 ± 21.86 vs. $39.68\pm26.61^\circ$), and calcium volume (1.42 ± 1.86 vs. 1.99 ± 2.87 mm³) for OCT and cryo-imaging respectively. There was good correlation between OCT and cryo for the previously mentioned parameters: $R=0.90$, 0.88 , and 0.76 , respectively ($p<0.01$). OCT underestimated calcium volume (3.11 ± 2.14 vs. 4.58 ± 3.39 mm³, $p<0.01$) in the 7 cases with deep calcifications as compared to cryo.

Conclusions: Intravascular OCT can accurately quantify the luminal depth, calcium volume and circumferential distribution of coronary plaque calcifications in humans.

